| | horo |
|--|------|

| Upland nurse crop | | |
|---------------------------------|------|---|
| Species per mix | 2 | |
| Seeds per square foot | 37 | |
| Native FQI NE Illinois | 0.00 | A consideration in choosing to use nurse crops is pollen production. The commonly used nurse crop grasses are wind pollinated. |
| Native FQI Illinois | 0.00 | Wind pollinated plants are the source of seasonal allergic reactions in many individuals. Some grasses are worse pollen sources |
| Native Mean C Value NE Illinois | 0.00 | than others. Among those known to cause allergic reactions include Annual Bluegrass, Sweet Vernal Grass, Bermuda Grass, Love |
| Native Mean C Value Illinois | 0.00 | Grasses, Kentucky Bluegrass, Orchard Grass, Paspalum, Perennial Rye Grass, Redtop, & Timothy. |
| Mean W Value NE Illlinois | 5.00 | |
| Mean W Value Illinois | 5.00 | |


| Graminoids, or grasses, sedges, rushes, and other grasslike plants. | | | | | | | | | | | | | | | | | | | | | |
|---|------------------------------------|----------|------------|------|-------|------|------|-------|------|---------------|---------|-----|--------|-------|---------------|----------|--------|-------------|------------|-------------|------------|
| ACRONYMS | S&W S&W wet Illinois IL wet IL wet | | | | | | | | | blo | om time | | | | | | | | | | |
| S&W ILL USDA vPlants scientific name | common name | family | type | C | index | acro | C | index | acro | height | color | MAN | M J | J A S | O N seeds/oz | seeds/lb | lb/ac | seeds/ acre | seeds/ftsq | % by weight | % by count |
| AVESAT AVESAT AVSA AVSA Avena sativa | Oats | Graminae | Ad A-Grass | * | 5 | UPL | * | 5 | UPL | 2.0-3.0 (2.5) | na | 9 | ⊕ ⊕ | | 938 | 15,000 | 32.000 | 480,000 | 11.02 | 86.49% | 29.72% |
| LOLMUL LOLMUL LOPEM2 LOMU Lolium multiflorum [L perenne m] | Italian Rye | Graminae | Ad A-Grass | * | 5 | UPL | * | 5 | UPL | 2.0-3.0 (2.5) | na | હ | e ee e | * * | 14,188 | 227,000 | 5.000 | 1,135,000 | 26.06 | 13.51% | 70.28% |
| v120812 | | | | | | | | | | | | _ | | | Graminoid tot | tals | 37.000 | 1,615,000 | 37 | 100.00% | 100.00% |
| | | | | 0.00 | 5.00 | | 0.00 | 5.00 | | | | | | | | | | | | | |

| Wetland nurse crop Species per mix | Knowingly designing plantings, or installing allergy-causing grasses in an urban context presents a severe liability exposure, on that presents more exposure than a one-day controlled burn. A clever lawyer or botanist with an allergic child living next to a planting with allergy causing nurse crops will sue someone, someday. It's is only a matter of time. You may be the first lawsuit think litigation. Think, "I say Errors & Omissions Insurance paid up?" Think responsibly. C.Y.A. The legal precedents for the exclusion of allergin producing landscape plants have been made. Be looking over your shoulder |
|--------------------------------------|---|
|--------------------------------------|---|

| Graminoids, or grasses, sedges, rushes, and other grasslike plants. | | | | | | | | | | | | | | | | | | | | |
|---|---------------|----------|------------|------|---------|---------|----------|--------|--------|---------------|-------|-----------|--------|--------------|-----------|--------|-------------|------------|-------------|------------|
| ACRONYMS | | | | S&W | S&W wet | S&W wet | Illinois | IL wet | IL wet | | | bloon | ı time | | | | | | | |
| S&W ILL USDA vPlants scientific name | common name | family | type | C | index | acro | C | index | acro | height | color | M A M J J | ASC | N seeds/oz | seeds/lb | lb/ac | seeds/ acre | seeds/ftsq | % by weight | % by count |
| AGRALA AGRALA AGGI2 AGGI2 Agrostis gigantea {A alba} | Red Top Grass | Graminae | Ad P-Grass | 0 | -3 | FACW | 0 | -3 | FACW | 2.5-3.5 (3.0) | na | ⊕ ⊕ | | 312,500 | 5,000,000 | 0.125 | 625,000 | 14.35 | 0.34% | 20.56% |
| AGRALP AGRALP AGST2 AGST2 Agrostis stolonifera (A alba palustri | s} Bent Grass | Graminae | Nt P-Grass | 10 | -5 | [OBL] | 8 | -3 | FACW | 1.0-2.0 (1.5) | na | ⊕ ⊕ | | 400,000 | 6,400,000 | 0.125 | 800,000 | 18.37 | 0.34% | 26.32% |
| AVESAT AVESAT AVSA AVSA Avena sativa | Oats | Graminae | Ad A-Grass | * | 5 | UPL | * | 5 | UPL | 2.0-3.0 (2.5) | na | ⊕ ⊕ | - | 938 | 15,000 | 32.000 | 480,000 | 11.02 | 85.91% | 15.79% |
| LOLMUL LOLMUL LOPEM2 LOMU Lolium multiflorum [L perenne m] | Italian Rye | Graminae | Ad A-Grass | * | 5 | UPL | * | 5 | UPL | 2.0-3.0 (2.5) | na | * * * | * * | 14,188 | 227,000 | 5.000 | 1,135,000 | 26.06 | 13.42% | 37.34% |
| v120812 | | | | | | | | | | | | | | Graminoid to | tals | 37.250 | 3,040,000 | 70 | 100.00% | 100.00% |
| | | | | 0.00 | 0.50 | | 0.00 | 1.00 | | | | | | | | | | | | |

| Savanna Nurse Crop | | |
|---------------------------------|------|--|
| Species per mix | 2 | |
| Seeds per square foot | 27 | Purely optional. If erosion or soil stability are a concern or they are your crutch, use one of the above nurse crops blended with you |
| Native FQI NE Illinois | 0.00 | native seeding. These mixes also act as efficient carriers & allow more efficient broadcasting of your seed mix. These nurse crop |
| Native FQI Illinois | 0.00 | mixes contain phytotoxic & highly allergenic species. Buyer Beware! |
| Native Mean C Value NE Illinois | 0.00 | |
| Native Mean C Value Illinois | 0.00 | |
| Mean W Value NE Illlinois | 5.00 | |
| Mean W Value Illinois | 5.00 | |

| Graminoids, or grasses, sedges, rushes, and other grasslike plants. | | | | | | | | | | | | | | | | | | | | | |
|---|----------------|----------|------------|------|---------|---------|----------|---------|--------|---------------|---------|---------|-------|-------|--------------|----------|--------|-------------|------------|-------------|------------|
| ACRONYMS | | | | S&W | S&W wet | S&W wet | Illinois | s IL we | IL wet | | | ŀ | oloom | ime | | | | | | | |
| S&W ILL USDA vPlants scientific name | common name | family | type | C | index | acro | C | index | acro | height | color ! | M A M J | J | A S O | N seeds/oz | seeds/lb | lb/ac | seeds/ acre | seeds/ftsq | % by weight | % by count |
| AVESAT AVESAT AVSA AVSA Avena sativa | Oats | Graminae | Ad A-Grass | * | 5 | UPL | * | 5 | UPL | 2.0-3.0 (2.5) | na | ⊕ ⊕ | | | 938 | 15,000 | 32.000 | 480,000 | 11.02 | 96.97% | 41.38% |
| FESOVI FESDUR FEOV FEOV FEOV Festuca ovina, F brevipila, F duriuscula | Sheep's Fescue | Graminae | Nt P-Grass | * | 5 | UPL | * | 5 | UPL | 1.0-2.0 (1.5) | na | • | • | | 42,500 | 680,000 | 1.000 | 680,000 | 15.61 | 3.03% | 58.62% |
| v120812 | | | | | | | | | | | | | | | Graminoid to | tals | 33.000 | 1,160,000 | 27 | 100.00% | 100.00% |
| | | | | 0.00 | 5.00 | | | 5.00 | | | | | | | | | | | | | |



Yellow is uncertain data